Foodborne viruses in Europe:

Web-based technologies for investigation of transnational outbreaks of viral gastroenteritis

Ben Lopman, for:

The Foodborne Viruses in Europe Consortium

M Koopmans, H Vennema, M Reacher, J Carrique-Mas, Y van Duynhoven, F-X Hanon and DWG Brown

Background

- Foodborne viral infections are increasingly recognised as a public health concern
- Norwalk-like viruses (NLV) are the most common cause of enteric disease
 - Large human reservoir
 - Stable outside the host
 - ➤ Small infectious dose (10-100 particles)
 - > Cause outbreaks among all age groups
- Multiple genetic variants of NLV co-circulate in the community

Background II

B B C NEWS


Front Page England Northern Ireland Scotland Wales **UK Politics** Business Sci/Tech Health Education **Entertainment Talking Point** In Depth

AudioVideo

You are in: UK: Scotland

Tuesday, 22 January, 2002, 20:38 GMT

World Virus leads to second hospital closure



The Victoria Infirmary remains closed

Background III





Health Officials Pinpoint Disease in Oscar Illness

EMAIL THIS ARTICLE

PRINTER FRIENDLY

BECOME A FOX FAN

Tuesday, March 19, 2002 Associated Press LOS ANGELES — A disease that spreads through food infected with sewage pollution likely is responsible for a

mystery illness that overcame at least 100 guests at a pre-Oscar ceremony earlier this month, public health officials said.


The March 2 event at the Regent Beverly Wilshire hotel in Beverly Hills honored scientific and technical achievement in cinema. About 500 people attended the dinner and awards presentation.

Days afterward, dozens of guests began complaining of vomiting, diarrhea and nausea. The illness lasted several days in most cases.

Burden of Norwalk-like virus gastroenteritis in US

Agents	Total Ilinesses	Foodborne Illnesses	% of all foodborne illness
All Bacteria	5,204,934	4,175,565	30.2
All Parasites	2,541,316	357,190	2.6
Norwalk-like viruses	23,000,000	9,200,000	66.6

Source: Mead et al, EID, 2000



Foodborne Viruses in Europe

University of Helsinki von Bonsdorff et al

Swedish Institute Infectious
Disease Control; Svensson et al

Statens Serum institute Bottiger et al

Robert Koch Institute Schreier et al

Central Public Health Institute Rome, Ruggeri et al

Objectives of the network

- 1) study the epidemiology of enteric viruses across Europe
- determine high risk foods & transmission routes
- 3) identify novel strains at the onset and trace their spread.
- 4) investigate mechanisms of emergence of these novel strains

Harmonisation of European Surveillance

Epidemiological Data

Virological Data





Development of web-accessible database

Harmonisation of European Surveillance

Epidemiological Data

Virological Data



Development of web-accessible database

Epidemiological Data I

- Harmonised clinical definitions
 - CASE of viral gastroenteritis
 - OUTBREAK of viral gastroenteritis
- Standard Outbreak Questionnaire
 - Etiology
 - Transmission
 - Setting
 - Case Information
 - Diagnostic Results
 - Food Vehicles

Epidemiological Data II

- Web-Based outbreak report form
- Active Server Pages technology
- Microsoft Access Database

http://www.eufoodborneviruses.net





Viral Gastroenteritis Outbreak Report Form



Fields with a * must be entered		
Reporting Institute :	Please select	*
Outbreak Reference:		*(your unique reference number)
Be sure to retain you outbreak reference. You w	ill need it to update the record.	
Reporter's Name:		*(First Last)
Today's date:		* dd/mm/yyyy
		submit ONLY virological data

Transmission

Mode of transmission:


Please select one


If foodborne, "PREPARED" takes precedence over "SERVED".
If person-ro-person "SERVED" takes precedence over "PREPARED".

Place where transmission of infection occurred:

Please select one

Was the outbreak the result of a point source exposure? (e.g. acafunction?)





Viral Gastroenteritis Outbreak UPDATE Form




Reporting Institute :	NL-National Institute for Public Health and the Environment 🔻 *		
Outbreak Number:	EP2002040	*(your unique reference number)	
Reporter's Name:	Yvonne van Duynhoven	*(First Last)	
Today's date:	14/03/2002	* dd/mm/yyyy	

Transmission

If foodborne, "PREPARED" takes precedence over "SERVED".

If person-to-person "SERVED" takes precedence over "PREPARED".

Mode of transmission:			
PERSON TO PERSON			
if "other", specify			
Place where transmission of infection occurred:			
Residential institution			
Specify:			
Was the outhreak the result of a point source evacure?/e.g.			





Viral Gastroenteritis Outbreak Search Page



Voormin	lotz wor	no artisa	CT 130/C	.++++++
Search	OV IE		$\simeq 1008$	шише.

Please select Search

Search by mode of transmission:

Please selectione Search

Search by setting:

Please selectione 🔻 Search



Click here to return to main page.

Short Outbreak report

Reporting Institute

NL-National Institute for Public Health and the Environment

Outbreak reference

EP2002040

Reporter's name

Yvonne van Duynhoven

Report date

14/03/2002

Mode of transmission/ setting

Mode of transmission Setting of outbreak

Details of setting

Was the outbreak the result of a point source exposure?

Date of point source exposure

PERSON TO PERSON Residential institution

nursing home

No

Aetiology

Was the organism identified?

Organism name:

Yes

Norwalk-like virus

Case information











Harmonisation of European Surveillance

Epidemiological Data

Virological Data





Harmonisation of methods



Development of web-accessible database

Virological Data: Sequencing of NLV






Virological Data II


Web-based genetic bank



Applied Maths, Ghent, Belgium




The databases are relational.





A group of linked outbreaks

Phylogenetic inference of emerging variant NLV IIb Hilversum:

Polymerase gene








A group of linked outbreaks

Phylogenetic inference of emerging variant NLV IIb Hilversum:



Summary

- We have identified linked outbreaks on an international scale
- Transnational outbreaks may be common
 - Introduce new genetic variant to an area
- The Foodborne Viruses in Europe group has developed a model for the detection of transnational outbreaks
 - Web accessible password protected database
 - Open to all participants
 - Links can be recognised from epidemiological OR virological characteristics of outbreaks

Conclusions

 Recognition of international outbreaks relies on both molecular typing and epidemiological data

 Preliminary results from our network show the value of standardised international databases for the recognition of transnational outbreaks

www.eufoodborneviruses.net

Acknowledgements

EU collaborators: David Brown, Jon Green, Mark Reacher, Marion Koopmans, Harry Vennema, Yvonne van Duynhoven, Wim van der Poel, Marc-Alain Widdowson, Lennart Svensson, Kjell-Olof Hedlund, Carl-Henrik von Bonsdorff, Leena Maunula, Kare Mölbak, Blenda Böttiger, Eckhart Schreier, Hans Gelderblom, Franco Ruggeri, Dario De Medici, Alicia Sánchez, Albert Bosch, Javier Buesa, Pierre Pottier, Eveline Kohli, Soizick LeGuyader, Juan-Carrique Mas, Elisabeth van Strien

EU contract QLK1-CT-1999-00594